



► Pr Catherine Bennetau-Pelissero  
Bordeaux Sciences Agro,  
Bordeaux France

## BIBLIOGRAPHIE

- [1] DiCanivenc-Lavier M-C, Bennetau-Pelissero C. Phytoestrogens and Health Effects. *Nutrients*. 2023;15(2):317.
- [2] Bensaada S, et al. Validation of ELISAs for Isoflavones and Enterolactone for Phytoestrogen Intake Assessment in the French Population. *Nutrients*. 2023;15(4):967.
- [3] Lee A et al. Endocrine disruptors on and in fruits and vegetables: Estimation of the potential exposure of the French population. *Food Chem*. 2022; 373(Pt B):131513.
- [4] NTP. Multigenerational reproductive study of genistein (Cas No. 446-72-0) in Sprague-Dawley rats (feed study). *Natl Toxicol Program Tech Rep Ser*. 2008;539:1-266.
- [5] NTP. Toxicology and carcinogenesis studies of genistein (Cas No. 446-72-0) in Sprague-Dawley rats (feed study). *Natl Toxicol Program Tech Rep Ser*. 2008;(545):1-240.
- [6] Bennetau-Pelissero C. Plant Proteins from Legumes. J.-M. Mérillon, K. G. Ramawat (eds.), *Bioactive Molecules in Food, Reference Series in Phytochemistry*. 2018;1-43
- [7] Bennetts HW, Underwood EJ, Shier FL. A specific breeding problem of sheep on subterranean clover pastures in Western Australia. *Aust Vet J*. 1946;22(1):2-12.
- [8] Farnsworth NR, Bingel AS, Cordell GA, et al. Potential value of plants as sources of new antifertility agents II. *J Pharm Sci*. 1975;64(5): 717-54.
- [9] Bennetau-Pelissero C. Natural Estrogenic Substances, Origins, and Effects. In J.-M. Mérillon, KG Ramawat (eds.), *Bioactive Molecules in Food, Reference Series in Phytochemistry*. 2018;1-70p.
- [10] Owens W, Ashby J, Odum J, et al. The OECD program to validate the rat uterotrophic bioassay. Phase 2: dietary phytoestrogen analyses. *Environ Health Perspect*. 2003;111(12):1559-67.
- [11] Perez-Rivero JJ, Martinez-Maya JJ, Perez-Martinez M, et al. Phytoestrogen treatment induces testis alterations in dogs. Potential use in population control. *Vet Res Commun*. 2009;33(1):87-95.
- [12] Golub MS, Hogrefe CE, Germann SL, et al. Neurobehavioral evaluation of rhesus monkey infants fed cow's milk formula, soy formula, or soy formula with added manganese. *Neurotoxicol Teratol*. 2005;27(4):615-27.
- [13] Patisaul HB. Endocrine disruption by dietary phyto-oestrogens: impact on dimorphic sexual systems and behaviours. *Proc Nutr Soc*. 2017;76(2):130-44.
- [14] Moon HJ, Seok JH, Kim SS, et al. Lactational coumestrol exposure increases ovarian apoptosis in adult rats. *Arch Toxicol*. 2009;83(6):601-8.
- [15] Stott P, Wight N. Female reproductive tract abnormalities in European hares (*Lepus europaeus*) in Australia. *J Wildl Dis*. 2004;40(4):696-703.
- [16] Sharpe RM, Martin B, Morris K, et al. Infant feeding with soy formula milk: effects on the testis and on blood testosterone levels in marmoset monkeys during the period of neonatal testicular activity. *Hum Reprod*. 2002;17(7):1692-703.
- [17] Index Mundi. <https://www.indexmundi.com/factbook/fields/age-structure> consulté le 30 Novembre 2023
- [18] Cunha GR, Li Y, Mei C, et al. Ontogeny of estrogen receptors in human male and female fetal reproductive tracts. *Differentiation*. 2021;118:107-31
- [19] Rochira V, Carani C. Estrogens, Male Reproduction and Beyond. In: Feingold KR, Anawalt B, Blackman MR, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; [Updated 2023 Mar 21]
- [20] Nagata C, Iwasa S, Shiraki M, et al. Associations among maternal soy intake, isoflavone levels in urine and blood samples, and maternal and umbilical hormone concentrations (Japan). *Cancer Causes Control*. 2006 Nov;17(9):1107-13.
- [21] North K, Golding J. A maternal vegetarian diet in pregnancy is associated with hypospadias. The ALSPAC Study Team. *Avon Longitudinal Study of Pregnancy and Childhood*. *BJU Int*. 2000; 85(1):107-13.
- [22] Akre O, Boyd HA, Ahlgren M, et al. Maternal and gestational risk factors for hypospadias. *Environ Health Perspect*. 2008;116(8):1071-6.
- [23] Michikawa T, Yamazaki S, Ono M, et al. Isoflavone Intake in Early Pregnancy and Hypospadias in the Japan Environment and Children's Study. *Urology*. 2019;124: 229-36.
- [24] Pasternski V, Acerini CL, Dunger DB, et al. Postnatal penile growth concurrent with mini-puberty predicts later sex-typed play behavior: Evidence for neurobehavioral effects of the postnatal androgen surge in typically developing boys. *Horm Behav*. 2015;69:98-105.
- [25] Afssa - Afssaps Sécurité et bénéfices des phyto-estrogènes apportés par l'alimentation - Recommandations Rapport d'expertise Mars 2005
- [26] McCarver G, Bhatia J, Chambers C, et al. NTP-CERHR Expert Panel Report on the Developmental Toxicity of Soy Infant Formula. *Birth Defects Research (Part B)*. 2011;92:421-68
- [27] Bennetau-Pelissero C, Sauvart P, Peltre G, et al. Phytoestrogènes du soja : Problèmes posés chez le nourrisson allergique au lait de vache et consommant les formules à base de soja. *Cah. Nutr. Diét*. 2004;39(1):24-33.
- [28] Harlid S, Adgent M, Jefferson WN, et al. Soy Formula and Epigenetic Modifications: Analysis of Vaginal Epithelial Cells from Infant Girls in the IFED Study. *Environ Health Perspect*. 2017;125(3):447-52.
- [29] Upson K, Harmon QE, Laughlin-Tommaso SK, et al. Soy-based Infant Formula Feeding and Heavy Menstrual Bleeding Among Young African American Women. *Epidemiology*. 2016;27(5):716-25.
- [30] Upson K, Adgent MA, Wegienka G, et al. Soy-based infant formula feeding and menstrual pain in a cohort of women aged 23-35 years. *Hum Reprod*. 2019;34(1):148-54.
- [31] Langton CR, Harmon QE, Upson K, et al. Soy-Based Infant Formula Feeding and Uterine Fibroid Development in a Prospective Ultrasound Study of Black/African-American Women. *Environ Health Perspect*. 2023;131(1):17006.
- [32] Upson K, Sathyanarayana S, Scholes D, et al. Early-life factors and endometriosis risk. *Fertil Steril*. 2015;104(4):964-971.e5.
- [33] Gilchrist JM, Moore MB, Andres A, et al. Ultrasonographic patterns of reproductive organs in infants fed soy formula: comparisons to infants fed breast milk and milk formula. *J Pediatr*. 2010;156(2):215-20.
- [34] Barthold JS, Manson J, Regan V, et al. Reproductive hormone levels in infants with cryptorchidism during postnatal activation of the pituitary-testicular axis. *J Urol*. 2004;172(4 Pt 2):1736-41
- [35] Kuiri-Hänninen T, Sankilampi U, Dunkel L. Activation of the hypothalamic-pituitary-gonadal axis in infancy: minipuberty. *Horm Res Paediatr*. 2014;82(2):73-80.
- [36] Faber KA, Hughes CL Jr. Dose-response characteristics of neonatal exposure to genistein on pituitary responsiveness to gonadotropin releasing hormone and volume of the sexually dimorphic nucleus of the preoptic area (SDN-POA) in postpubertal castrated female rats. *Reprod Toxicol*. 1993;7(1):35-9.
- [37] Adgent MA, Daniels JL, Edwards LJ, et al. Early-life soy exposure and gender-role play behavior in children. *Environ Health Perspect*. 2011;119(12):1811-6.
- [38] Kim J, Kim S, Huh K, et al. High serum isoflavone concentrations are associated with the risk of precocious puberty in Korean girls. *Clin Endocrinol (Oxf)*. 2011;75(6): 831-5.
- [39] Cheng G, Remer T, Prinz-Langenohl R, et al. Relation of isoflavones and fiber intake in childhood to the timing of puberty. *Am J Clin Nutr*. 2010;92(3):556-64.
- [40] Messina M, Rogero MM, Fisberg M, et al. Health impact of childhood and adolescent soy consumption. *Nutr Rev*. 2017;75(7):500-15.
- [41] Wada K, Nakamura K, Masue T, et al. Soy intake and urinary sex hormone levels in preschool Japanese children. *Am J Epidemiol*. 2011;173(9):998-1003.
- [42] Kunisue T, Tanabe S, Isobe T et al. Profiles of Phytoestrogens in Human Urine from Several Asian Countries. *J. Agric. Food Chem*. 2010;58:9838-9846.
- [43] Liu Z, et al. Intake of soy foods and soy isoflavones by rural adult women in China. *J. Asia Pac J Clin Nutr*. 2004;13(2):204-9.
- [44] Hooper L, Ryder JJ, Kurzer MS, et al. Effects of soy protein and isoflavones on circulating hormone concentrations in pre- and post-menopausal women: a systematic review and meta-analysis. *Human Reprod Update*. 2009;15(4):423-40.
- [45] Andrews MA, Schliep KC, Wactawski-Wende J, et al. Dietary factors and luteal phase deficiency in healthy eumenorrheic women. *Hum Reprod*. 2015;30(8):1942-51.
- [46] Jacobsen BK, Jaceldo-Siegl K, Knutsen SF, et al. Soy isoflavone intake and the likelihood of ever becoming a mother: the Adventist Health Study-2. *Int J Women's Health*. 2014;6:377-84.
- [47] Chandrareddy A, Munejyirici-Delale O, McFarlane SI, et al. Adverse effects of phytoestrogens on reproductive health: a report of three cases. *Complement Ther Clin Pract*. 2008;14(2):132-5.
- [48] Imai H, Nishikawa H, Suzuki A, et al. Secondary Hypogonadism due to Excessive Ingestion of Isoflavone in a Man. *Intern Med*. 2022;61(19):2899-903.
- [49] Pitsouni E, Grigoriadis T, Douskos A, et al. Efficacy of vaginal therapies alternative to vaginal estrogens on sexual function and orgasm of menopausal women: A systematic review and meta-analysis of randomized controlled trials. *Eur J Obstet Gynecol Reprod Biol*. 2018;229:45-56.
- [50] Liu J, Yuan F, Gao J, et al. Oral isoflavone supplementation on endometrial thickness: a meta-analysis of randomized placebo-controlled trials. *Oncotarget*. 2016;7(14):17369-79.
- [51] Bennetau-Pelissero C. Phyto-estrogènes et cancers du sein *Phyto Europ*. 136:8-16
- [52] Mirzaee F, Fakari FR, Babakhanian M, et al. The Effectiveness of Herbal Medicines on Cyclic Mastalgia: A Systematic Review on Meta-analysis. *Rev Bras Ginecol Obstet*. 2022;44(10):972-985.